



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Patient Blood Management: success and potential in the future

Spahn, Donat R

DOI: <https://doi.org/10.1097/SLA.0000000000001787>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-128879>

Journal Article

Published Version

Originally published at:

Spahn, Donat R (2016). Patient Blood Management: success and potential in the future. *Annals of Surgery*, 264(2):212-213.

DOI: <https://doi.org/10.1097/SLA.0000000000001787>

Patient Blood Management

Success and Potential in the Future

Donat R. Spahn, MD, FRCA

Preoperative anemia,¹ exaggerated blood loss,² and allogeneic red blood cell (RBC) transfusions³ are individual and intertwined factors that adversely affect patient outcome. They are modifiable and represent the 3 pillars of patient blood management (PBM)—a comprehensive and interdisciplinary concept that aims at pre-emptively reducing the need for RBC transfusions to improve patient safety and outcome.⁴ The 3 pillars of PBM are: treatment of preoperative anemia, reduction of perioperative RBC loss, and optimization of the physiologic anemia tolerance including the use of restrictive transfusion triggers.

Meybohm et al, in this issue of the Journal, show that the implementation of PBM in all surgical disciplines in 4 German University Hospitals is safe and reduces RBC transfusions by 17% and acute renal failure by 30%.⁵ It is important to stress that this represents real-life data because all surgical patients were included without any exclusion criteria. Safety, the primary outcome, was assessed as the composite outcome of in-hospital mortality, myocardial infarction, ischemic stroke, acute renal failure, pneumonia, or sepsis. It is highly interesting (Tables 3 and 5 of the paper⁵) that RBC transfusions and the primary outcome progressively decreased the longer PBM had been implemented. I hypothesize that the full benefit of PBM had not yet been achieved 1 to 2 years after PBM implementation because the RBC transfusion rate also progressively decreased over several years after the implementation of a PBM program in Western Australia.⁶

Given the multitude of measures of a comprehensive PBM program, it is not astonishing that some centers did not yet fully succeed in the implementation of certain PBM measures (Fig. 1B of the paper⁵). Particularly well implemented was the emphasis on surgical hemostasis and the point-of-care diagnostic for coagulation monitoring. In contrast, relatively little progress was achieved so far in the extended preoperative diagnostics and treatment of anemia. Consequently, the anemia rate at admission remained high at a staggering 35% and admission hemoglobin concentration remained stable. With unchanged discharge hemoglobin, but reduced RBC transfusions, the perioperative RBC loss must have gone down. The exact reason has not been formally analyzed in this study, but likely represents a reduced surgical blood loss due to improved surgical hemostasis and coagulation monitoring and treatment.

The yet unchanged preoperative anemia rate represents a promising field for future research and unmet PBM potential. Preoperative anemia affects approximately 30% of patients and is associated with increased perioperative mortality (20%–40%), increased morbidity (30%), more RBC transfusions, and prolonged length of hospital stay.^{1,7} Anemia treatment is efficacious in orthopedic,^{8,9} cardiac,¹⁰ and abdominal surgery,^{11,12} not only in reducing RBC transfusions but also in reducing acute kidney injury, infections, and length of hospital stay.^{8–12} Why then is preoperative

From the Institute of Anesthesiology, University and University Hospital of Zurich, Zurich, Switzerland.

Received for publication April 12, 2016; accepted April 13, 2016.

Reprints: Donat R. Spahn, MD, FRCA Professor and Chairman Institute of Anesthesiology Section Head Medical Anesthesiology - Intensive Care Medicine - OR-Management University of Zurich and University Hospital Zurich Raemistrasse 100 CH-8091 Zurich, Switzerland. E-mail: donat.spahn@usz.ch.

Conflict of interest statement (past 5 years): Donat R. Spahn—Dr Spahn's academic department is/has been receiving grant support from the Swiss National Science Foundation, Berne, Switzerland, the Ministry of Health (Gesundheitsdirektion) of the Canton of Zurich, Switzerland for Highly Specialized Medicine, the Swiss Society of Anesthesiology and Reanimation (SGAR), Berne, Switzerland, the Swiss Foundation for Anesthesia Research, Zurich, Switzerland, Bundesprogramm Chancengleichheit, Berne, Switzerland, CSL Behring, Berne, Switzerland, Vifor SA, Villars-sur-Glâne, Switzerland.

Dr Spahn was the chair of the ABC Faculty and is the co-chair of the ABC-Trauma Faculty, managed by Physicians World Europe GmbH, Mannheim, Germany and sponsored by unrestricted educational grants from Novo Nordisk Health Care AG, Zurich, Switzerland, CSL Behring GmbH, Marburg, Germany and LFB Biomédicaments, Courtaboeuf Cedex, France.

In the past 5 years, Dr Spahn has received honoraria or travel support for consulting or lecturing from: Abbott AG, Baar, Switzerland, AMGEN GmbH, Munich, Germany, AstraZeneca AG, Zug, Switzerland, Baxter AG, Volketswil, Switzerland, Baxter S.p.A., Roma, Italy, Bayer, Zürich, Switzerland and Berlin, Germany, B. Braun Melsungen AG, Melsungen, Germany, Boehringer Ingelheim (Schweiz) GmbH, Basel, Switzerland, Bristol-Myers-Squibb, Rueil-Malmaison Cedex, France and Baar, Switzerland, CSL Behring GmbH, Hattersheim am Main, Germany and Berne, Switzerland, Curacyte AG, Munich, Germany, Daiichi Sankyo (Schweiz) AG, Thalwil, Switzerland, Ethicon Biosurgery, Sommerville, New Jersey, USA, Fresenius SE, Bad Homburg v.d.H., Germany, Galenica AG, Bern, Switzerland (including Vifor SA, Villars-sur-Glâne, Switzerland), GlaxoSmithKline GmbH & Co. KG, Hamburg, Germany, Janssen-Cilag, Baar, Switzerland and Beersse, Belgium, LFB Biomédicaments, Courtaboeuf Cedex, France, Merck Sharp & Dohme AG, Luzern, Switzerland, Novo Nordisk A/S, Bagsvård, Denmark, Octapharma AG, Lachen, Switzerland, Organon AG, Pfäffikon/SZ, Switzerland, PAION Deutschland GmbH, Aachen, Germany, Pharmacosmos A/S, Holbaek, Denmark, Photonics Healthcare B.V., Utrecht, Netherlands, ratiopharm Arzneimittel Vertriebs-GmbH, Vienna, Austria, Roche, Reinach, Switzerland, Schering-Plough International, Inc., Kenilworth, New Jersey, USA, Tem International GmbH, Munich, Germany, Verum Diagnostica GmbH, Munich, Germany, Vifor Pharma, Munich, Germany, Vienna, Austria and St. Gallen, Switzerland.

Copyright © 2016 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 0003-4932/14/26105-0821

DOI: 10.1097/SLA.0000000000001787

anemia treatment not better implemented yet in clinical practice? The most important reasons are that many physicians:

1. insufficiently know the definition of anemia, that is, a hemoglobin <12 g/dL for women and a hemoglobin <13 g/dL for men,
2. do not fully recognize the perioperative burden of even mild anemia, that is, a hemoglobin of 10 to 12 g/dL for women and a hemoglobin of 10 to 13 g/dL for men,
3. report patients only late to anesthesiologists and therefore there seems to be little time for anemia detection and treatment,
4. perceive anemia treatment to be costly and requiring up-front investment.

In addition, systematic preoperative anemia treatment is indeed logistically demanding in large hospitals.⁶ Nevertheless, we are to invest efforts into a better anemia treatment in the future. The Duke Perioperative Enhancement Team recently delineated in detail how to develop and implement a preoperative anemia clinic successfully.¹³ In addition to early referral, “intelligent scheduling systems” are conceivable. Such systems would allow the scheduling of elective patients for operations with an expected blood loss of more than say 500 mL or a probability of RBC transfusion greater than 10% only if preoperative anemia had been investigated and successfully treated. Preoperative anemia treatment programs thus may finally become standard clinical practice.^{6,14}

Unfortunately, the authors did not analyze costs. This is another important aspect for the acceptance of a new treatment paradigm not only for physicians but also for hospital administrators and third-party payers. The authors are encouraged to use their data set for a formal cost analysis because the introduction of PBM programs and PBM measures have been shown to be cost-saving in the past in smaller studies.^{13,15–18}

I congratulate the authors of this study and all the participating physicians for their tremendous effort in implementing PBM systematically in their hospitals. They have proven safety and benefits of this concept. The full potential of PBM, however, is certainly larger, and the study clearly highlights areas of future points of action, first of which may be the implementation of structures for systematic preoperative anemia treatment.

REFERENCES

1. Musallam KM, Tamim HM, Richards T, et al. Preoperative anaemia and postoperative outcomes in non-cardiac surgery: a retrospective cohort study. *Lancet*. 2011;378:1396–1407.
2. Ranucci M, Baryshnikova E, Castelvechio S, et al. Major bleeding, transfusions, and anemia: the deadly triad of cardiac surgery. *Ann Thorac Surg*. 2013;96:478–485.
3. Spahn DR, Goodnough LT. Alternatives to blood transfusion. *Lancet*. 2013;381:1855–1865.
4. Spahn DR, Moch H, Hofmann A, et al. Patient blood management: the pragmatic solution for the problems with blood transfusions. *Anesthesiology*. 2008;109:951–953.
5. Meybohm P, Herrmann E, Steinbicker AU, et al. Patient blood management is associated with a substantial reduction of red blood cell utilization and safe for patient's outcome: a prospective, multicenter cohort study with a noninferiority design. *Ann Surg*. 2016;264:203–211.
6. Leahy MF, Roberts H, Mukhtar SA, et al. A pragmatic approach to embedding patient blood management in a tertiary hospital. *Transfusion*. 2014;54:1133–1145.
7. Baron DM, Hochrieser H, Posch M, et al. Preoperative anaemia is associated with poor clinical outcome in non-cardiac surgery patients. *Br J Anaesth*. 2014;113:416–423.
8. Munoz M, Gomez-Ramirez S, Cuenca J, et al. Very-short-term perioperative intravenous iron administration and postoperative outcome in major orthopedic surgery: a pooled analysis of observational data from 2547 patients. *Transfusion*. 2014;54:289–299.
9. Rineau E, Chaudet A, Chassier C, et al. Implementing a blood management protocol during the entire perioperative period allows a reduction in transfusion rate in major orthopedic surgery: a before-after study. *Transfusion*. 2016;56:673–681.
10. Yoo YC, Shim JK, Kim JC, et al. Effect of single recombinant human erythropoietin injection on transfusion requirements in preoperatively anemic patients undergoing valvular heart surgery. *Anesthesiology*. 2011;115:929–937.
11. Froessler B, Palm P, Weber I, et al. The important role for intravenous iron in perioperative patient blood management in major abdominal surgery: a randomized controlled trial. *Ann Surg*. 2016.
12. Calleja JL, Delgado S, Del Val A, et al. Ferric carboxymaltose reduces transfusions and hospital stay in patients with colon cancer and anemia. *Int J Colorectal Dis*. 2016;31:543–551.
13. Guinn NR, Guercio JR, Hopkins TJ, et al. How do we develop and implement a preoperative anemia clinic designed to improve perioperative outcomes and reduce cost? *Transfusion*. 2016;56:297–303.
14. Spahn DR, Zacharowski K. Non-treatment of preoperative anaemia is substandard clinical practice. *Br J Anaesth*. 2015;115:1–3.
15. Gross I, Seifert B, Hofmann A, et al. Patient blood management in cardiac surgery results in fewer transfusions and better outcome. *Transfusion*. 2015;55:1075–1081.
16. Mehra T, Seifert B, Bravo-Reiter S, et al. Implementation of a patient blood management monitoring and feedback program significantly reduces transfusions and costs. *Transfusion*. 2015;55:2807–2815.
17. Pearse BL, Smith I, Faulke D, et al. Protocol guided bleeding management improves cardiac surgery patient outcomes. *Vox Sang*. 2015;109:267–279.
18. Spahn DR, Theusinger OM, Hofmann A. Patient blood management is a win-win: a wake-up call. *Br J Anaesth*. 2012;108:889–892.